



# Lecture 7. Adaptive Online Convex Optimization

Advanced Optimization (Fall 2024)

**Peng Zhao**

`zhaop@lamda.nju.edu.cn`

Nanjing University

# Outline

- Motivation
  - Minimax results
  - Beyond the worst-case analysis
  - Problem-dependent consideration
- Small-Loss Bounds
  - Small-loss bound for PEA
  - Self-confident Tuning
  - Small-loss bound for OCO

# Part 1. Motivation

- Minimax Results
- Beyond the worst-case analysis
- Problem-dependent guarantees

# General Regret Analysis for OMD

## Online Mirror Descent

$$\mathbf{x}_{t+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} \left\{ \eta_t \langle \mathbf{x}, \nabla f_t(\mathbf{x}_t) \rangle + \mathcal{D}_\psi(\mathbf{x}, \mathbf{x}_t) \right\}$$

**Theorem 4** (General Regret Bound for OMD). Assume  $\psi$  is  $\lambda$ -strongly convex w.r.t.  $\|\cdot\|$  and  $\eta_t = \eta, \forall t \in [T]$ . Then, for all  $\mathbf{u} \in \mathcal{X}$ , the following regret bound holds

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq \frac{\mathcal{D}_\psi(\mathbf{u}, \mathbf{x}_1)}{\eta} + \frac{\eta}{\lambda} \sum_{t=1}^T \|\nabla f_t(\mathbf{x}_t)\|_\star^2 - \frac{1}{\eta} \sum_{t=1}^T \mathcal{D}_\psi(\mathbf{x}_{t+1}, \mathbf{x}_t)$$

# Online Mirror Descent


- Our previous mentioned algorithms can **all be covered**

*minimax  
optimal*

Algo.	OMD/proximal form	$\psi(\cdot)$	$\eta_t$	Regret <sub>T</sub>
OGD for convex	$\mathbf{x}_{t+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} \eta_t \langle \mathbf{x}, \nabla f_t(\mathbf{x}_t) \rangle + \frac{1}{2} \ \mathbf{x} - \mathbf{x}_t\ _2^2$	$\ \mathbf{x}\ _2^2$	$\frac{1}{\sqrt{t}}$	$\mathcal{O}(\sqrt{T})$
OGD for strongly c.	$\mathbf{x}_{t+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} \eta_t \langle \mathbf{x}, \nabla f_t(\mathbf{x}_t) \rangle + \frac{1}{2} \ \mathbf{x} - \mathbf{x}_t\ _2^2$	$\ \mathbf{x}\ _2^2$	$\frac{1}{\sigma t}$	$\mathcal{O}(\frac{1}{\sigma} \log T)$
ONS for exp-concave	$\mathbf{x}_{t+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} \eta_t \langle \mathbf{x}, \nabla f_t(\mathbf{x}_t) \rangle + \frac{1}{2} \ \mathbf{x} - \mathbf{x}_t\ _{A_t}^2$	$\ \mathbf{x}\ _{A_t}^2$	$\frac{1}{\gamma}$	$\mathcal{O}(\frac{d}{\gamma} \log T)$
Hedge for PEA	$\mathbf{x}_{t+1} = \arg \min_{\mathbf{x} \in \Delta_N} \eta_t \langle \mathbf{x}, \nabla f_t(\mathbf{x}_t) \rangle + \text{KL}(\mathbf{x} \ \mathbf{x}_t)$	$\sum_{i=1}^N x_i \log x_i$	$\sqrt{\frac{\ln N}{T}}$	$\mathcal{O}(\sqrt{T \log N})$

# Beyond the Worst-Case Analysis


- All above regret guarantees hold against the worst case
  - Matching the *minimax optimality*
  - The environment is *fully adversarial*
- However, in practice:
  - We are not always interested in the *worst-case scenario*
  - Environments can exhibit *specific patterns*: gradual change, periodicity...



We are after *problem-dependent* guarantees.

# Beyond the Worst-Case Analysis

- Beyond the worst-case analysis, achieving more adaptive results.
  - (1) *adaptivity*: achieving better guarantees in easy problem instances;
  - (2) *robustness*: maintaining the same worst-case guarantee.



[Slides from Dylan Foster, [Adaptive Online Learning](#) @NIPS'15 workshop]

# Prediction with Expert Advice

- Recall the PEA setup

At each round  $t = 1, 2, \dots$

- (1) the player first picks a weight  $\mathbf{p}_t$  from a simplex  $\Delta_N$ ;
- (2) and simultaneously environments pick a loss vector  $\ell_t \in \mathbb{R}^N$ ;
- (3) the player suffers loss  $f_t(\mathbf{p}_t) \triangleq \langle \mathbf{p}_t, \ell_t \rangle$ , observes  $\ell_t$  and updates the model.

- Performance measure: *regret*

$$\text{Regret}_T \triangleq \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - \min_{i \in [N]} \sum_{t=1}^T \ell_{t,i}$$

*benchmark the performance with respect to the **best expert***



# Part 2. Small-loss for PEA

- Refined Analysis for Hedge
- Self-confident Tuning

# Hedge: Regret Bound

**Theorem 1.** Suppose that  $\forall t \in [T]$  and  $i \in [N], 0 \leq \ell_{t,i} \leq 1$ , then Hedge with learning rate  $\eta$  guarantees

$$\text{Regret}_T \leq \frac{\ln N}{\eta} + \eta T = \mathcal{O}(\sqrt{T \log N}),$$

*minimax optimal*

where the last equality is by setting  $\eta$  optimally as  $\sqrt{(\ln N)/T}$ .

- What if there exists an *excellent* expert? i.e.,  $L_{T,i} \ll T$  holds for some  $i \in [N]$ .
- Goal: can we achieve a “*small-loss*” bound? something like  $\mathcal{O}(\sqrt{L_{T,i^*} \log N})$ .

# Small-Loss Bounds for PEA

**Theorem 2.** Suppose that  $\forall t \in [T]$  and  $i \in [N]$ ,  $0 \leq \ell_{t,i} \leq 1$ , then Hedge with learning rate  $\eta \in (0, 1)$  guarantees

$$\sum_{t=1}^T \langle \mathbf{p}_t, \boldsymbol{\ell}_t \rangle - \min_{i \in [N]} \sum_{t=1}^T \ell_{t,i} \leq \frac{1}{1-\eta} \left( \frac{\ln N}{\eta} + \eta L_{T,i^*} \right),$$

by setting  $\eta = \min \left\{ \frac{1}{2}, \sqrt{\frac{\ln N}{L_{T,i^*}}} \right\}$ , we have the following small-loss regret bound:

$$\text{Regret}_T = \mathcal{O} \left( \sqrt{L_{T,i^*} \log N} + \log N \right).$$

- (1) **adaptivity**: when  $L_{T,i^*} = \mathcal{O}(1)$ , the regret bound is  $\mathcal{O}(\log N)$ , which is independent of  $T$ !
- (2) **robustness**: when  $L_{T,i^*} = \mathcal{O}(T)$ , it can recover the *minimax*  $\mathcal{O}(\sqrt{T \log N})$  guarantee.

# Improved Analysis for Small-Loss Bound

*Proof.*

$$\sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \frac{\ln N}{\eta} + \eta \sum_{t=1}^T \sum_{i=1}^N p_{t,i} \ell_{t,i}^2$$

- For previous worst-case analysis, we simply utilize  $\ell_{t,i} \leq 1$ :

$$\eta \sum_{t=1}^T \sum_{i=1}^N p_{t,i} \ell_{t,i}^2 \leq \eta T$$

- To get a small-loss bound, we **improve** the analysis to be:

$$\eta \sum_{t=1}^T \sum_{i=1}^N p_{t,i} \ell_{t,i}^2 \leq \eta \sum_{t=1}^T \sum_{i=1}^N p_{t,i} \ell_{t,i} = \eta \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle$$

$$\Rightarrow \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \frac{\ln N}{\eta} + \eta \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle$$

# Improved Analysis for Small-Loss Bound

*Proof.*  $\Rightarrow \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \frac{\ln N}{\eta} + \eta \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle$

$$(1 - \eta) \left( \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \right) \leq \frac{\ln N}{\eta} + \eta L_{T,i^*} \quad (\text{rearrange})$$

$$\Rightarrow \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \frac{1}{1 - \eta} \left( \frac{\ln N}{\eta} + \eta L_{T,i^*} \right)$$

**Lemma 1.** Let  $a, b > 0$  and  $x_0 > 0$  be three positive values. Suppose that  $L \leq ax + \frac{b}{x}$  holds for any  $x \in (0, x_0]$ . Then, by taking  $x^* = \min\{\sqrt{b/a}, x_0\}$ , we have  $L \leq 2\sqrt{ab} + \frac{2b}{x_0}$ .

# Improved Analysis for Small-Loss Bound

*Proof.*  $\implies \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \frac{1}{1-\eta} \left( \frac{\ln N}{\eta} + \eta L_{T,i^*} \right)$

**Lemma 1.** Let  $a, b > 0$  and  $x_0 > 0$  be three positive values. Suppose that  $L \leq ax + \frac{b}{x}$  holds for any  $x \in (0, x_0]$ . Then, by taking  $x^* = \min\{\sqrt{b/a}, x_0\}$ , we have  $L \leq 2\sqrt{ab} + \frac{2b}{x_0}$ .

*Proof.* Suppose  $\sqrt{b/a} \leq x_0$ , then  $x^* = \sqrt{b/a}$  and we have  $L \leq ax^* + \frac{b}{x^*} = 2\sqrt{ab}$ . Otherwise,  $x^* = x_0$  and we have  $L \leq ax^* + \frac{b}{x^*} = ax_0 + \frac{b}{x_0}$ . Notice that in latter case  $x_0 \leq \sqrt{b/a}$  holds, which implies  $ax_0 \leq \frac{b}{x_0}$  and hence  $ax_0 + \frac{b}{x_0} \leq \frac{2b}{x_0}$ . Combining two cases ends the proof.  $\square$

# Improved Analysis for Small-Loss Bound

*Proof.*  $\implies \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \frac{1}{1-\eta} \left( \frac{\ln N}{\eta} + \eta L_{T,i^*} \right)$

**Lemma 1.** Let  $a, b > 0$  and  $x_0 > 0$  be three positive values. Suppose that  $L \leq ax + \frac{b}{x}$  holds for any  $x \in (0, x_0]$ . Then, by taking  $x^* = \min\{\sqrt{b/a}, x_0\}$ , we have  $L \leq 2\sqrt{ab} + \frac{2b}{x_0}$ .


Therefore, we get an  $\mathcal{O}(\sqrt{L_{T,i^*} \log N} + \log N)$  small-loss regret

by setting the learning rate optimally as  $\eta^* = \min\left\{\frac{1}{2}, \sqrt{\frac{\ln N}{L_{T,i^*}}}\right\}$ .  $\square$


# Learning Rate Tuning Issue

Therefore, we get an  $\mathcal{O}(\sqrt{L_{T,i^*} \log N} + \log N)$  small-loss regret

by setting the learning rate optimally as  $\eta^* = \min \left\{ \frac{1}{2}, \sqrt{\frac{\ln N}{L_{T,i^*}}} \right\}$ .



However, this online algorithm is not legitimate, due to the requirement of using  $L_{T,i^*}$  (the cumulative loss of the best expert) as the input.



*Fortunately, we can remedy it by the **self-confident tuning** framework.*



# Self-confident Tuning Framework

- Recall the OGD algorithm for convex function:

$$\mathbf{x}_{t+1} = \Pi_{\mathcal{X}} [\mathbf{x}_t - \eta_t \nabla f_t(\mathbf{x}_t)]$$

which enjoys the following regret bound

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \min_{\mathbf{x} \in \mathcal{X}} \sum_{t=1}^T f_t(\mathbf{x}) \leq \frac{D^2}{\eta} + \eta G^2 T.$$

We can set  $\eta = \frac{D}{G\sqrt{\textcolor{red}{T}}}$  to obtain an  $\mathcal{O}(\sqrt{T})$  regret bound.

**Question:** can we remove the dependence of  $\textcolor{red}{T}$  when tuning the step size?

$\Rightarrow$  A natural guess is to set  $\eta_t = \frac{D}{G\sqrt{\textcolor{blue}{t}}}$ .

# Self-confident Tuning Framework

- **Self-confident tuning**: utilize the available empirical quantities to approximate the unknown ones.

⇒ use  $\eta_t = \frac{D}{G\sqrt{t}}$  to approximate  $\eta^* = \frac{D}{G\sqrt{T}}$ , ensuring the same bound (in order).

**Theorem 3.** Suppose the diameter of non-empty closed convex set  $\mathcal{X}$  is  $D$  and  $\|\nabla f_t(\mathbf{x})\| \leq G$  for any  $\mathbf{x} \in \mathcal{X}$ . Then OGD with step size tuning  $\eta_t = \frac{D}{G\sqrt{t}}$  ensures the following regret bound:

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq \frac{3}{2}GD\sqrt{T}.$$

# Self-confident Tuning Framework

**Theorem 3.** Suppose the diameter of non-empty closed convex set  $\mathcal{X}$  is  $D$  and  $\|\nabla f_t(\mathbf{x})\| \leq G$  for any  $\mathbf{x} \in \mathcal{X}$ . Then OGD with step size tuning  $\eta_t = \frac{D}{G\sqrt{t}}$  ensures the following regret bound:

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq \frac{3}{2}GD\sqrt{T}.$$

**Proof.**

$$\begin{aligned} \sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{u}) &\leq \frac{1}{2} \sum_{t=1}^T \left( \frac{1}{\eta_t} - \frac{1}{\eta_{t-1}} \right) \|\mathbf{u} - \mathbf{x}_t\|_2^2 + \sum_{t=1}^T \eta_t \|\nabla f_t(\mathbf{x}_t)\|_2^2 \\ &\leq \frac{D^2}{2} \sum_{t=1}^T \left( \frac{1}{\eta_t} - \frac{1}{\eta_{t-1}} \right) + \frac{G^2}{2} \sum_{t=1}^T \eta_t \\ &= \frac{D^2}{2\eta_T} + \frac{GD}{2} \sum_{t=1}^T \frac{1}{\sqrt{t}} \quad \left( \frac{1}{\eta_0} = 0 \right) \\ &\leq \frac{GD\sqrt{T}}{2} + GD\sqrt{T} = \frac{3}{2}GD\sqrt{T} \quad \left( \sum_{t=1}^T \frac{1}{\sqrt{t}} \leq 2\sqrt{T} \right) \end{aligned}$$

# Self-confident Tuning Framework

- Consider the small-loss bound for PEA problem.

Achieving small loss bound  $\mathcal{O}(\sqrt{L_{T,i^*} \log N} + \log N)$  with  $\eta = \min \left\{ \frac{1}{2}, \sqrt{\frac{\ln N}{L_{T,i^*}}} \right\}$ .

**Goal:** tuning  $\eta$  without the knowledge of  $L_{T,i^*}$

Deploying self-confident tuning: how can we empirically approximate  $L_{T,i^*}$ ?

$$L_{T,i} \triangleq \sum_{t=1}^T \ell_{t,i}, \quad i^* = \arg \min_{i \in [N]} L_{T,i}$$

$$L_{t,i} \triangleq \sum_{s=1}^t \ell_{s,i}, \quad i_t^* = \arg \min_{i \in [N]} L_{t,i}$$

$\Rightarrow$  **Key challenge:** index  $i^*$  and index sequence  $\{i_t^*\}_{t=1}^T$  can be highly different

# Self-confident Tuning Framework

- Consider the small-loss bound for PEA problem.

Achieving small loss bound  $\mathcal{O}(\sqrt{L_{T,i^*} \log N} + \log N)$  with  $\eta = \min \left\{ \frac{1}{2}, \sqrt{\frac{\ln N}{L_{T,i^*}}} \right\}$ .

We need to dive into the regret analysis.

$$\sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \frac{\ln N}{\eta} + \eta \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle$$

$$\Rightarrow \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq 2\sqrt{(\ln N) \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle}$$

by setting  $\eta = \sqrt{\frac{\ln N}{\sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle}}$ .

$\Leftrightarrow$  Denoted by  $\tilde{L}_T = \sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle$   
we obtain  $\tilde{L}_T - L_{T,i^*} \leq 2\sqrt{(\ln N) \tilde{L}_T}$

**Lemma.** For  $x, y, a \in \mathbb{R}_+$  that satisfy  $x - y \leq \sqrt{ax}$ , it implies  $x - y \leq \sqrt{ay} + a$ .

$\Rightarrow \tilde{L}_T - L_{T,i^*} \leq 2\sqrt{(\ln N) L_{T,i^*}} + 4 \ln N$   
by resolving  $\tilde{L}_T$ .

# Self-confident Tuning Framework

- Consider the small-loss bound for PEA problem.

Achieving small loss bound  $\mathcal{O}(\sqrt{L_{T,i^*} \log N} + \log N)$  with  $\eta = \min \left\{ \frac{1}{2}, \sqrt{\frac{\ln N}{L_{T,i^*}}} \right\}$ .

More specifically, setting  $\eta = \sqrt{\frac{\ln N}{\tilde{L}_T}}$ , yields

$$\tilde{L}_T - L_{T,i^*} \leq \frac{\ln N}{\eta} + \eta \tilde{L}_T \implies \tilde{L}_T - L_{T,i^*} \leq 2\sqrt{(\ln N) \tilde{L}_T} \implies \tilde{L}_T - L_{T,i^*} \leq \mathcal{O}(\sqrt{(\log N) L_{T,i^*}} + \log N)$$

While  $\tilde{L}_T$  cannot be obtained ahead of time, a *natural* empirical approximation is:

$$\eta_t = \sqrt{\frac{\ln N}{\tilde{L}_t}}, \quad \text{where } \tilde{L}_t = \sum_{s=1}^t \langle \mathbf{p}_s, \ell_s \rangle \quad p_{t+1,i} \propto \exp(-\eta_t L_{t,i}), \quad \forall i \in [N]$$

# Self-confident Tuning Framework

**Theorem 4.** Suppose that  $\forall t \in [T]$  and  $i \in [N], 0 \leq \ell_{t,i} \leq 1$ , then Hedge with adaptive learning rate  $\eta_t = \sqrt{\frac{\ln N}{\tilde{L}_t + 1}}$  guarantees

$$\begin{aligned}\text{Regret}_T &\leq 6\sqrt{(\mathbf{L}_{T,i^*} + 1) \ln N} + 36 \ln N \\ &= \mathcal{O}\left(\sqrt{\mathbf{L}_{T,i^*} \log N} + \log N\right),\end{aligned}$$

where  $\tilde{L}_t = \sum_{s=1}^t \langle \mathbf{p}_s, \ell_s \rangle$  is cumulative loss the learner suffered at time  $t$ .

# Proof

*Proof.* We again use ‘potential-based’ proof here, where the **potential** is defined as

$$\begin{aligned}\Phi_t(\eta) &\triangleq \frac{1}{\eta} \ln \left( \sum_{i=1}^N \exp(-\eta L_{t,i}) \right) \\ \Phi_t(\eta_{t-1}) - \Phi_{t-1}(\eta_{t-1}) &= \frac{1}{\eta_{t-1}} \ln \left( \frac{\sum_{i=1}^N \exp(-\eta_{t-1} L_{t,i})}{\sum_{i=1}^N \exp(-\eta_{t-1} L_{t-1,i})} \right) \\ &= \frac{1}{\eta_{t-1}} \ln \left( \sum_{i=1}^N \left( \frac{\exp(-\eta_{t-1} L_{t-1,i})}{\sum_{i=1}^N \exp(-\eta_{t-1} L_{t-1,i})} \exp(-\eta_{t-1} \ell_{t,i}) \right) \right) \\ &= \frac{1}{\eta_{t-1}} \ln \left( \sum_{i=1}^N p_{t,i} \exp(-\eta_{t-1} \ell_{t,i}) \right) \quad (\text{update rule of } \mathbf{p}_t) \\ &\quad (p_{t,i} \propto \exp(-\eta_{t-1} L_{t-1,i}), \forall i \in [N])\end{aligned}$$



# Proof

*Proof.*

$$\begin{aligned}\Phi_t(\eta_{t-1}) - \Phi_{t-1}(\eta_{t-1}) &= \frac{1}{\eta_{t-1}} \ln \left( \sum_{i=1}^N p_{t,i} \exp(-\eta_{t-1} \ell_{t,i}) \right) \\ &\leq \frac{1}{\eta_{t-1}} \ln \left( \sum_{i=1}^N p_{t,i} (1 - \eta_{t-1} \ell_{t,i} + \eta_{t-1}^2 \ell_{t,i}^2) \right) \quad (\forall x \geq 0, e^{-x} \leq 1 - x + x^2) \\ &= \frac{1}{\eta_{t-1}} \ln \left( 1 - \eta_{t-1} \langle \mathbf{p}_t, \boldsymbol{\ell}_t \rangle + \eta_{t-1}^2 \sum_{i=1}^N p_{t,i} \ell_{t,i}^2 \right) \\ &\leq -\langle \mathbf{p}_t, \boldsymbol{\ell}_t \rangle + \eta_{t-1} \sum_{i=1}^N p_{t,i} \ell_{t,i}^2 \quad (\ln(1+x) \leq x)\end{aligned}$$

# Proof

$$\Phi_t(\eta) \triangleq \frac{1}{\eta} \ln \left( \sum_{i=1}^N \exp(-\eta L_{t,i}) \right)$$

*Proof.* 
$$\Phi_t(\eta_{t-1}) - \Phi_{t-1}(\eta_{t-1}) \leq -\langle \mathbf{p}_t, \boldsymbol{\ell}_t \rangle + \eta_{t-1} \sum_{i=1}^N p_{t,i} \ell_{t,i}^2$$

$$\Rightarrow \quad \langle \mathbf{p}_t, \boldsymbol{\ell}_t \rangle \leq \Phi_{t-1}(\eta_{t-1}) - \Phi_t(\eta_{t-1}) + \eta_{t-1} \sum_{i=1}^N p_{t,i} \ell_{t,i}^2$$

$$\begin{aligned} \sum_{t=1}^T \langle \mathbf{p}_t, \boldsymbol{\ell}_t \rangle &\leq \Phi_0(\eta_0) - \Phi_T(\eta_{T-1}) + \sum_{t=1}^T \left( \eta_{t-1} \sum_{i=1}^N p_{t,i} \ell_{t,i}^2 \right) + \sum_{t=1}^T \left( \Phi_t(\eta_t) - \Phi_t(\eta_{t-1}) \right) \quad (\text{telescoping}) \\ &\leq \frac{\ln N}{\eta_{T-1}} - \frac{1}{\eta_{T-1}} \ln(\exp(-\eta_{T-1} L_{T,i^*})) + \sum_{t=1}^T \eta_{t-1} \sum_{i=1}^N p_{t,i} \ell_{t,i}^2 + \sum_{t=1}^T (\Phi_t(\eta_t) - \Phi_t(\eta_{t-1})) \quad (\eta_0 \geq \eta_{T-1}) \\ &= \sqrt{(\tilde{L}_{T-1} + 1) \ln N} + L_{T,i^*} + \sum_{t=1}^T \eta_{t-1} \langle \mathbf{p}_t, \boldsymbol{\ell}_t \rangle + \sum_{t=1}^T (\Phi_t(\eta_t) - \Phi_t(\eta_{t-1})) \quad (\ell_{t,i} \leq 1) \end{aligned}$$

# Proof

$$\Phi_t(\eta) \triangleq \frac{1}{\eta} \ln \left( \sum_{i=1}^N \exp(-\eta L_{t,i}) \right)$$

**Proof.** 
$$\sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle \leq \sqrt{(\tilde{L}_{T-1} + 1) \ln N} + L_{T,i^*} + \sum_{t=1}^T \eta_{t-1} \langle \mathbf{p}_t, \ell_t \rangle + \sum_{t=1}^T \left( \Phi_t(\eta_t) - \Phi_t(\eta_{t-1}) \right)$$

To bound  $\sum_{t=1}^T \left( \Phi_t(\eta_t) - \Phi_t(\eta_{t-1}) \right)$ , we prove that  $\Phi_t(\eta)$  is increasing w.r.t.  $\eta$ :

$$\begin{aligned} \eta^2 \Phi'_t(\eta) &= \eta^2 \left( -\frac{1}{\eta^2} \ln \left( \frac{1}{N} \sum_{i=1}^N \exp(-\eta L_{t,i}) \right) - \frac{1}{\eta} \frac{\sum_{i=1}^N L_{t,i} \exp(-\eta L_{t,i})}{\sum_{i=1}^N \exp(-\eta L_{t,i})} \right) \\ &= \ln N - \sum_{i=1}^N p_{t+1,i}^\eta \left( \ln \left( \sum_{j=1}^N \exp(-\eta L_{t,j}) \right) + \eta L_{t,i} \right) \quad (p_{t+1,i}^\eta \propto \exp(-\eta L_{t,i})) \\ &= \ln N - \sum_{i=1}^N p_{t+1,i}^\eta \ln \left( \frac{\sum_{j=1}^N \exp(-\eta L_{t,j})}{\exp(-\eta L_{t,i})} \right) \\ &= \ln N - \sum_{i=1}^N p_{t+1,i}^\eta \ln \frac{1}{p_{t+1,i}^\eta} \geq 0 \quad \Rightarrow \sum_{t=1}^T \left( \Phi_t(\eta_t) - \Phi_t(\eta_{t-1}) \right) \leq 0 \quad (\eta_t \leq \eta_{t-1}) \end{aligned}$$

# Proof

*Proof.* From the potential-based proof, we already know that

$$\sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \sqrt{(\tilde{L}_{T-1} + 1) \ln N} + \sum_{t=1}^T \eta_{t-1} \langle \mathbf{p}_t, \ell_t \rangle$$

$$\leq \sqrt{(\tilde{L}_{T-1} + 1) \ln N} + \sqrt{\ln N} \cdot \sum_{t=1}^T \frac{\langle \mathbf{p}_t, \ell_t \rangle}{\sqrt{\sum_{s=1}^{t-1} \langle \mathbf{p}_s, \ell_s \rangle + 1}} \quad \begin{array}{l} (\eta_{t-1} = \sqrt{\frac{\ln N}{\tilde{L}_{t-1} + 1}}) \\ (\tilde{L}_{t-1} = \sum_{s=1}^{t-1} \langle \mathbf{p}_s, \ell_s \rangle) \end{array}$$

How to bound this term?

$\Rightarrow$  A common structure to handle.

# Self-confident Tuning Lemma

**Lemma 2.** *Let  $a_1, a_2, \dots, a_T$  be non-negative real numbers. Then*

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^t a_s}} \leq 2 \sqrt{1 + \sum_{t=1}^T a_t}$$

**Lemma 3.** *Let  $a_1, a_2, \dots, a_T$  be non-negative real numbers. Then*

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^{t-1} a_s}} \leq 4 \sqrt{1 + \sum_{t=1}^T a_t + \max_{t \in [T]} a_t}$$

The two lemmas are useful for analyzing algorithms with self-confident tuning.

# Proof

**Lemma 2.** Let  $a_1, a_2, \dots, a_T$  be non-negative real numbers. Then

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^t a_s}} \leq 2 \sqrt{1 + \sum_{t=1}^T a_t}$$

*Proof.*

$$\frac{1}{2}x \leq 1 - \sqrt{1 - x}, \forall x \in [0, 1]$$

Let  $a_0 \triangleq 1$ , by set  $x = a_t / \sum_{s=0}^t a_s$ :

$$\frac{a_t}{2 \sum_{s=0}^t a_s} \leq 1 - \sqrt{1 - \frac{a_t}{\sum_{s=0}^t a_s}}$$

# Proof

*Proof.*

$$\frac{a_t}{2 \sum_{s=0}^t a_s} \leq 1 - \sqrt{1 - \frac{a_t}{\sum_{s=0}^t a_s}}$$

$$\Rightarrow \frac{a_t}{2\sqrt{\sum_{s=0}^t a_s}} \leq \sqrt{\sum_{s=0}^t a_s} - \sqrt{\sum_{s=0}^t a_s - \sum_{s=0}^{t-1} a_s}$$

By telescoping from  $t = 1$  to  $T$ :

$$\sum_{t=1}^T \left( \frac{a_t}{2\sqrt{1 + \sum_{s=1}^t a_s}} \right) \leq \sqrt{\sum_{s=0}^T a_s} - \sqrt{\sum_{s=0}^1 a_s - \sum_{s=0}^0 a_s} \leq \sqrt{1 + \sum_{t=1}^T a_t} \quad \square$$

# Proof

**Lemma 3.** Let  $a_1, a_2, \dots, a_T$  be non-negative real numbers. Then

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^{t-1} a_s}} \leq 4 \sqrt{1 + \sum_{t=1}^T a_t + \max_{t \in [T]} a_t}$$

*Proof.* We define that  $\max_{t \in [T]} a_t = B$ .

• **Case 1.** If  $\sum_{t=1}^T a_t \leq B$ :

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^{t-1} a_s}} \leq \sum_{t=1}^T a_t \leq B, \text{ Lemma 2 is obviously satisfied.}$$



# Proof

**Lemma 3.** Let  $a_1, a_2, \dots, a_T$  be non-negative real numbers. Then

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^{t-1} a_s}} \leq 4 \sqrt{1 + \sum_{t=1}^T a_t + \max_{t \in [T]} a_t}$$

**Proof.** We define that  $\max_{t \in [T]} a_t = B$ .

- **Case 2.** If  $\sum_{t=1}^T a_t \geq B$ , we define  $t_0 \triangleq \min \left\{ t : \sum_{s=1}^{t-1} a_s \geq B \right\}$ :

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^{t-1} a_s}} \leq B + \sum_{t=t_0}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^{t-1} a_s}} \leq B + \sum_{t=t_0}^T \frac{a_t}{\sqrt{1 + \frac{\sum_{s=1}^{t-1} a_s + a_t}{2}}}$$

$(\frac{x+y}{2} \leq x \text{ for } x \geq y)$

# Proof

**Lemma 3.** Let  $a_1, a_2, \dots, a_T$  be non-negative real numbers. Then

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^{t-1} a_s}} \leq 4 \sqrt{1 + \sum_{t=1}^T a_t + \max_{t \in [T]} a_t}$$

**Proof.** We define that  $\max_{t \in [T]} a_t = B$ .

- **Case 2.** If  $\sum_{t=1}^T a_t \geq B$ , we define  $t_0 \triangleq \min \left\{ t : \sum_{s=1}^{t-1} x_s \geq B \right\}$ :

$$B + \sum_{t=t_0}^T \frac{a_t}{\sqrt{1 + \frac{\sum_{s=1}^{t-1} a_s + a_t}{2}}} \leq B + \sum_{t=t_0}^T \frac{2a_t}{\sqrt{1 + \sum_{s=1}^t a_s}} \stackrel{(\text{Lemma 1})}{\leq} B + 4 \sqrt{1 + \sum_{t=1}^T a_t} \quad \square$$

# Small-Loss bound for PEA: Proof

*Proof.* From previous potential-based proof, we already know that

$$\sum_{t=1}^T \langle \mathbf{p}_t, \ell_t \rangle - L_{T,i^*} \leq \sqrt{(\tilde{L}_{T-1} + 1) \ln N} + \sqrt{\ln N} \cdot \sum_{t=1}^T \frac{\langle \mathbf{p}_t, \ell_t \rangle}{\sqrt{\sum_{s=1}^{t-1} \langle \mathbf{p}_s, \ell_s \rangle + 1}}$$

**Lemma 3.** Let  $a_1, a_2, \dots, a_T$  be non-negative real numbers. Then

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^{t-1} a_s}} \leq 4 \sqrt{1 + \sum_{t=1}^T a_t + \max_{t \in [T]} a_t}$$

$$\begin{aligned} \Rightarrow \quad \tilde{L}_T - L_{T,i^*} &\leq \sqrt{(\tilde{L}_{T-1} + 1) \ln N} + \sqrt{\ln N} \cdot \left( 4 \sqrt{1 + \tilde{L}_T + 1} \right) \quad (\ell_i \leq 1, \forall i \in [N]) \\ &\leq 5 \sqrt{(\tilde{L}_T + 1) \ln N} + \sqrt{\ln N} \end{aligned}$$

# Small-Loss bound for PEA: Proof

*Proof.*  $\implies \tilde{L}_T - L_{T,i^*} \leq 5\sqrt{(\tilde{L}_T + 1) \ln N} + \sqrt{\ln N}$

By the lemma, let  $x = \tilde{L}_T + 1, y = L_{T,i^*} + 1$ :

**Lemma.** For  $x, y, a \in \mathbb{R}_+$  that satisfy  $x - y \leq \sqrt{ax}$ , it implies  $x - y \leq \sqrt{ay} + a$ .

$$(\tilde{L}_T + 1) - (L_{T,i^*} + 1) \leq 6\sqrt{(\tilde{L}_T + 1) \ln N}$$

This implies that

$$(\tilde{L}_T + 1) - (L_{T,i^*} + 1) \leq 6\sqrt{(L_{T,i^*} + 1) \ln N} + 36 \ln N$$

$$\implies \tilde{L}_T - L_{T,i^*} = \mathcal{O}\left(\sqrt{L_{T,i^*} \log N} + \log N\right). \quad \square$$

# Part 3. Small-loss for OCO

- Small-loss quantity for OCO
- Small-loss OGD and self-confident tuning

# Small-loss PEA to OCO

- We have obtained a PEA algorithm with small-loss bound.

**Theorem 4.** Suppose that  $\forall t \in [T]$  and  $i \in [N]$ ,  $0 \leq \ell_{t,i} \leq 1$ , then Hedge with adaptive learning rate  $\eta_t = \sqrt{\frac{\ln N}{\tilde{L}_t + 1}}$  guarantees

$$\begin{aligned}\text{Regret}_T &\leq 6\sqrt{(L_{T,i^*} + 1) \ln N} + 36 \ln N \\ &= \mathcal{O}\left(\sqrt{L_{T,i^*} \log N} + \log N\right),\end{aligned}$$

where  $\tilde{L}_t = \sum_{s=1}^t \langle \mathbf{p}_s, \ell_s \rangle$  is cumulative loss the learner suffered at time  $t$ .

- Can we further extend the result to more **general OCO** setting?

# Small Loss in General OCO Setting

**Definition 4** (Small Loss). The small-loss quantity of the OCO problem (online function  $f_t : \mathcal{X} \mapsto \mathbb{R}$ ) is defined as

$$F_T = \min_{\mathbf{x} \in \mathcal{X}} \sum_{t=1}^T f_t(\mathbf{x})$$

- By taking  $f_t(\mathbf{x}) = \langle \mathbf{x}, \ell_t \rangle$  and  $\mathcal{X} = \Delta_N$ , we recover the definition of the small-loss quantity of PEA problem:

$$F_T = \min_{\mathbf{x} \in \Delta_N} \sum_{t=1}^T \langle \mathbf{x}, \ell_t \rangle = \sum_{t=1}^T \ell_{t,i^*} = L_{T,i^*}$$

# Small Loss in General OCO Setting

**Definition 4** (Small Loss). The small-loss quantity of the OCO problem (online function  $f_t : \mathcal{X} \mapsto \mathbb{R}$ ) is defined as

$$F_T = \min_{\mathbf{x} \in \mathcal{X}} \sum_{t=1}^T f_t(\mathbf{x})$$

A possible target regret bound:

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq \mathcal{O}(\sqrt{1 + F_T}).$$



# Self-bounding Property

- We require the following *self-bounding property* to ensure the small-loss bound for general OCO.

**Lemma 4** (Self-bounding Property). For an *L-smooth* function  $f : \mathbb{R}^d \mapsto \mathbb{R}$  with  $\mathbf{x}^* \in \arg \min_{\mathbf{v} \in \mathbb{R}^d} f(\mathbf{v})$ , we have that

$$\|\nabla f(\mathbf{x})\|_2 \leq \sqrt{2L(f(\mathbf{x}) - f(\mathbf{x}^*))}, \quad \forall \mathbf{x} \in \mathcal{X}.$$

**Corollary 1.** For an *L-smooth* and *non-negative* function  $f : \mathbb{R}^d \mapsto \mathbb{R}$ , we have that

$$\|\nabla f(\mathbf{x})\|_2 \leq \sqrt{2Lf(\mathbf{x})}, \quad \forall \mathbf{x} \in \mathcal{X}.$$

# Achieving Small-Loss Bound

- We show that under the *self-bounding condition*, OGD can yield the desired small-loss regret bound.

$$\mathbf{x}_{t+1} = \Pi_{\mathbf{x} \in \mathcal{X}} [\mathbf{x}_t - \eta_t \nabla f_t(\mathbf{x}_t)]$$

**Theorem 6** (Small-loss Bound). Assume that  $f_t$  is  $L$ -smooth and non-negative for all  $t \in [T]$ , when setting  $\eta_t = \frac{D}{\sqrt{1+\tilde{G}_t}}$ , the regret of OGD to any comparator  $\mathbf{u} \in \mathcal{X}$  is bounded as

$$\text{Regret}_T = \sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq \mathcal{O} \left( \sqrt{1 + F_T} \right)$$

where  $\tilde{G}_t = \sum_{s=1}^t \|\nabla f_s(\mathbf{x}_s)\|_2^2$  is the empirical estimator of cumulative gradient  $G_T$ .

# Proof

**Proof.** 
$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq \sum_{t=1}^T \eta_t \|\nabla f_t(\mathbf{x}_t)\|_2^2 + \sum_{t=1}^T \frac{1}{2\eta_t} \left( \|\mathbf{u} - \mathbf{x}_t\|_2^2 - \|\mathbf{u} - \mathbf{x}_{t+1}\|_2^2 \right)$$

$$\sum_{t=1}^T \eta_t \|\nabla f_t(\mathbf{x}_t)\|_2^2 = D \underbrace{\sum_{t=2}^T \frac{\|\nabla f_t(\mathbf{x}_t)\|_2^2}{\sqrt{1 + \tilde{G}_t}}}_{(\eta_1 \triangleq 1)} + G^2 \leq 2D \sqrt{1 + \sum_{t=1}^T \|\nabla f_t(\mathbf{x}_t)\|_2^2} + G^2$$

$(\tilde{G}_t = \sum_{s=1}^t \|\nabla f_s(\mathbf{x}_s)\|_2^2)$

**Lemma 2.** *Let  $a_1, a_2, \dots, a_T$  be non-negative real numbers. Then*

$$\sum_{t=1}^T \frac{a_t}{\sqrt{1 + \sum_{s=1}^t a_s}} \leq 2 \sqrt{1 + \sum_{t=1}^T a_t}$$

# Proof

**Proof.** 
$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq \sum_{t=1}^T \eta_t \|\nabla f_t(\mathbf{x}_t)\|_2^2 + \sum_{t=1}^T \frac{1}{2\eta_t} \left( \|\mathbf{u} - \mathbf{x}_t\|_2^2 - \|\mathbf{u} - \mathbf{x}_{t+1}\|_2^2 \right)$$

$$\sum_{t=1}^T \eta_t \|\nabla f_t(\mathbf{x}_t)\|_2^2 = D \sum_{t=2}^T \frac{\|\nabla f_t(\mathbf{x}_t)\|_2^2}{\sqrt{1 + \tilde{G}_t}} + G^2 \leq 2D \sqrt{1 + \sum_{t=1}^T \|\nabla f_t(\mathbf{x}_t)\|_2^2} + G^2$$

$(\eta_1 \triangleq 1)$   $(\tilde{G}_t = \sum_{s=1}^t \|\nabla f_s(\mathbf{x}_s)\|_2^2)$

$$\leq 2D \sqrt{1 + 2L \sum_{t=1}^T f_t(\mathbf{x}_t)} + G^2$$

(self-bounding property)

# Proof

**Proof.** 
$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq \sum_{t=1}^T \eta_t \|\nabla f_t(\mathbf{x}_t)\|_2^2 + \sum_{t=1}^T \frac{1}{2\eta_t} \left( \|\mathbf{u} - \mathbf{x}_t\|_2^2 - \|\mathbf{u} - \mathbf{x}_{t+1}\|_2^2 \right)$$

$$\sum_{t=1}^T \eta_t \|\nabla f_t(\mathbf{x}_t)\|_2^2 \leq 2D \sqrt{1 + 2L \sum_{t=1}^T f_t(\mathbf{x}_t)} + G^2$$

$$\sum_{t=1}^T \frac{1}{2\eta_t} \left( \|\mathbf{u} - \mathbf{x}_t\|_2^2 - \|\mathbf{u} - \mathbf{x}_{t+1}\|_2^2 \right) \leq \frac{D}{2} \sqrt{1 + 2L \sum_{t=1}^T f_t(\mathbf{x}_t)} + \frac{D}{2}$$

$$\Rightarrow \text{Regret}_T = \sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq 3D \sqrt{1 + 2L \sum_{t=1}^T f_t(\mathbf{x}_t)} + G^2$$

# Proof

**Proof.** 
$$\text{Regret}_T = \sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) \leq 3D \sqrt{1 + 2L \sum_{t=1}^T f_t(\mathbf{x}_t) + G^2}$$

Remember how we solve a similar problem in PEA:

## Small-Loss bound for PEA: Proof

**Proof.**  $\Rightarrow \tilde{L}_T - L_{T,i^*} \leq \sqrt{(\tilde{L}_T + 1) \ln N} + 4\sqrt{1 + \tilde{L}_T} + 1$

Then we solve above inequality. Let  $x \triangleq \tilde{L}_T + 1$ :

$$x - (\sqrt{\ln N} + 4)\sqrt{x} \leq L_{T,i^*} + 2 \quad \Rightarrow \quad \left( \sqrt{x} - \frac{\sqrt{\ln N} + 4}{2} \right)^2 \leq L_{T,i^*} + 2 + \left( \frac{\sqrt{\ln N} + 4}{2} \right)^2$$

This implies that

$$\sqrt{\tilde{L}_T} - 1 \leq \sqrt{L_{T,i^*} + 2 + \left( \frac{\sqrt{\ln N} + 4}{2} \right)^2} + \frac{\sqrt{\ln N} + 4}{2}$$

$$\Rightarrow \tilde{L}_T \leq 3 \ln N + L_{T,i^*} + 8\sqrt{(L_{T,i^*} + 1) \ln N} = \mathcal{O}(\sqrt{L_{T,i^*} \log N} + \log N). \quad (\text{squaring both sides}) \quad \square$$

Advanced Optimization (Fall 2024)

Lecture 7. Adaptive Online Convex Optimization

39

$$\Rightarrow \text{Regret}_T = \sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{u}) = \mathcal{O} \left( D \sqrt{L \sum_{t=1}^T f_t(\mathbf{u}) + 1 + G^2} \right). \quad \square$$

# Proof of Self-bounding Property

**Lemma 4** (Self-bounding Property). For an *L-smooth* function  $f : \mathbb{R}^d \mapsto \mathbb{R}$  with  $\mathbf{x}^* \in \arg \min_{\mathbf{v} \in \mathbb{R}^d} f(\mathbf{v})$ , we have that

$$\|\nabla f(\mathbf{x})\|_2 \leq \sqrt{2L(f(\mathbf{x}) - f(\mathbf{x}^*))}, \quad \forall \mathbf{x} \in \mathcal{X}.$$

**Proof.** By smoothness over the entire  $\mathbb{R}^d$  space, we have for any  $\mathbf{x}, \boldsymbol{\delta} \in \mathbb{R}^d$

$$f(\mathbf{x} + \boldsymbol{\delta}) \leq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \boldsymbol{\delta} \rangle + \frac{L}{2} \|\boldsymbol{\delta}\|_2^2.$$

Choosing  $\boldsymbol{\delta} = -\frac{\nabla f(\mathbf{x})}{L}$  gives  $f\left(\mathbf{x} - \frac{\nabla f(\mathbf{x})}{L}\right) \leq f(\mathbf{x}) - \frac{\|\nabla f(\mathbf{x})\|_2^2}{2L}$ .

(actually one-step improvement lemma)

Notice that  $f(\mathbf{x}^*) \leq f\left(\mathbf{x} - \frac{\nabla f(\mathbf{x})}{L}\right)$  by definition, which implies

$$f(\mathbf{x}^*) \leq f\left(\mathbf{x} - \frac{\nabla f(\mathbf{x})}{L}\right) \leq f(\mathbf{x}) - \frac{\|\nabla f(\mathbf{x})\|_2^2}{2L}.$$


Rearranging the above terms finishes the proof. □

# Several Remarks

- Remark 1: about the non-negative assumption  
When the online functions are non-negative, it is possible to redefine the small-loss quantity by incorporating each-round minimal function value.
- Remark 2: about the smoothness assumption  
Smoothness is necessary to obtain small-loss regret bound by the first-order method (can be proved by the online-to-batch conversion and existing lower bounds for deterministic optimization).
- Remark 3: take care of the way dealing with variance term  
In OGD here we use Lemma 1, while in Hedge for PEA we use Lemma 2.



# Summary



Q & A

*Thanks!*